
New England Governors/Eastern Canadian Premiers

Acid Rain Action Plan
Action Item 4 : Forest Mapping Project

Mapping Forest Sensitivity to Atmospheric Acid Deposition

2006-2007 Annual Report

Prepared by

The Forest Mapping Group

For

The Committee on the Environment of
The Conference of New England Governors and Eastern Canadian Premiers

Summary

The Forest Mapping Work Group has undertaken to map the sensitivity of the entire New England Governor and Eastern Canadian premier (NEG/ECP) jurisdictions' forests to atmospheric sulfur and nitrogen deposition loadings. Unprecedented for level of detail and size of area studied, this comprehensive project is the first scientific large-scale study of forest sensitivity to sulfur and nitrogen deposition in northeastern North America.

Critical Load

The term 'critical load' implies a tipping point, or threshold. Most generally, critical load may be defined as the maximum load that a system can tolerate before failing. As applied to environmental issues, however, critical load usually refers to exposure to pollutants; a critical load is an estimate of the level of exposure to one or more pollutants below which no harmful effects are known to occur to specified elements within an environment.

The common definition of critical loads agreed upon by the Forest Mapping Group is the definition originally developed by the United Nations Economic Commission for Europe in 1988:

A critical load is a quantitative estimate of the exposure to one or more pollutants below which significant harmful effects on specific sensitive elements of the environment do not occur according to present knowledge (Nilsson and Grennfelt, 1988).

The use of critical loads within the context of air quality management is premised on the notion that the effectiveness air quality policy is reflected in ecosystem impacts. The critical load concept is uniquely well suited toward informing air quality policy because its receptor-based approach takes into account both the spatial and topographical variables of atmospheric deposition.

As it applies to the atmospheric deposition of acid forming compounds then, a critical load is that level of exposure to sulfur and nitrogen compounds below which no harmful effects are known to occur within a specified environment (or ecosystem).

A unified critical load map for NEG/ECP region is now being completed; for this map, critical loads have been calculated for Northeastern North American forest ecosystem.

The approach used to identify critical loads for sulfur and nitrogen in the region's forest ecosystem is an ecological assessment based on an overall (steady-state) ecosystem budget for nutrient cations of calcium (Ca^{2+}), magnesium (Mg^{2+}), and potassium (K^+). This budget exists within a dynamic system of nutrient inputs, exports, and recycling.

In its simplest terms, the inputs to the nutrient budget for the Maine forest ecosystem include the addition of the nutrients Ca, Mg, and K through atmospheric deposition; acid forming compounds of sulfur (S) and nitrogen (N) are also introduced through

deposition. Additional inputs of Ca, Mg, and K are occur through the chemical weathering of the bedrock and soils.

Nutrient losses or exports from the system occur as a result of chemical reactions within the root zone which may render a portion of nutrients unavailable for plant nutrition, and through soil leaching in response to the presence of acids. Additional losses or exports occur as a result of forest fires and through the harvesting of trees from the forest.

Nutrient recycling occurs throughout the lifecycle of the trees in the forest through the shedding of leaves and/or needles and through the decay of vegetative and woody debris on the forest floor.

The overall ecosystem budget is based upon the relative values of the inputs to and exports from the system. A condition where nutrient input values exceed exports suggests that an equilibrium state of biologic capacity exists for that ecosystem. Conversely, a condition where nutrient exports exceed imports suggests a net nutrient deficit and increasing soil acidification; conditions ultimately unsustainable for that ecosystem over the long term.

The critical load map developed for the NEG/ECP region is derived on the basis of steady-state or static models. Consequently the map reflects conditions of nutrient balance rather than absolute measures of soil acidity/fertility. Nevertheless one might observe that where a negative nutrient imbalance is small, forest health problems and growth decline may not yet be evident; in those locations where the imbalance is significant, the impacts on forest health are likely to be observable today.

Critical load approaches offer air quality and natural resource managers a powerful tool with which to identify ecosystems at risk and to tailor management strategies to address specific resource issues.

Results

The Critical Load mapping performed for the Region indicates that up to 61% of forested land within certain of the region's jurisdictions have been characterized as 'sensitive', and thus may be experiencing a net nutrient deficit and increasing soil acidification. Soil mineral nutrient depletion has been linked to a wide variety of forest health problems, including reduced growth rates and increased mortality.

Figure 1. Forest areas sensitive to acid deposition in the New England states and Eastern Canadian provinces.
 (Please note discussion of data variability on page 7)




Table 1. Forest soil critical loads by NEG/ECP jurisdictions and their exceedances.
 Values are area-weighted. The terrestrial mapped area covers 162,674 km² in the U.S. and 547,424 km² in Canada.

Jurisdiction	Critical Load		Exceedance		
	Median	5 th	Median	95 th	Area mapped as exceeded
		percentile		percentile exceedance	
(eq ha ⁻¹ y ⁻¹)					(%)
Maine	1280	340	-420	660	35.8
New Hampshire	1350	440	-520	600	17.6
Vermont	1600	390	-390	930	29.9
Rhode Island	1130	230	70	1260	51.6
Massachusetts	1770	540	-420	1250	29.1
Connecticut	2290	1330	-790	-10	4.4
Total New England states	1590	360	-470	730	29.3
Newfoundland	519	227	16	343	52.3
Nova Scotia	692	353	81	538	61.2
Prince-Edward-Island	2212	876	-1549	-190	3.3
New Brunswick	960	488	-215	298	28.2
Quebec	934	420	-175	532	31.6
Total Eastern Canadian provinces	946	423	-220	444	37.6

Table 2. Comparison of statewide average metrics for elements of the forest sensitivity assessment in the New England states to show the relative importance of forest management activities compared with acid deposition.

Jurisdiction	S+N Deposition eq ha ⁻¹ y ⁻¹	S+N Range ¹	Base Cation Leaching eq ha ⁻¹ y ⁻¹	Base Cation Harvesting eq ha ⁻¹ y ⁻¹	Base Cation Weathering eq ha ⁻¹ y ⁻¹	WX SD ²	Critical Load eq ha ⁻¹ y ⁻¹
Maine	680	1320	530	410	2440	1870	1280
New Hampshire	900	2050	810	260	1800	0430	1350
Vermont	1010	2100	890	230	2500	2090	1600
Rhode Island	1290	870	1360	330	2200	330	1130
Massachusetts	1310	1510	1400	210	2270	1240	1770
Connecticut	1350	1180	1450	170	2680	1150	2290
Newfoundland	528	416	601	47	383	394	616
Nova Scotia	739	381	638	69	445	507	633
Prince-Edward- Island	637	97	1386	71	1221	767	1922
New Brunswick	681	902	984	132	885	625	1051
Quebec	770	980	190	60	740	1690	930

Highlights

- Forest areas sensitive to acid deposition reach 253,685 km² (35.7%) of the mapped area for the NEG/ECP jurisdictions (Figure 1, Table 1).
- In the Eastern Canadian provinces, the most sensitive forest areas occur in Southern Quebec, especially in the Lower-Laurentides located at the north of the St. Lawrence River, in Southeastern Nova Scotia, and in Southern Newfoundland.
- In New England, the most sensitive forest areas occur in the mountain ranges and coastal areas where soils are poor and weathering rates low, and where there is greater demand for nutrients due to more intensive harvesting (Table 2).
- The greatest forest sensitivity areas correspond to locations where acid deposition rates are high (south and west areas), and where critical loads are low (varies with geology and harvesting rate – Table 2).
- Direct comparisons between jurisdictions should be made with caution since they were mapped at different resolutions and accuracy depending on data availability. Variations in data availability and scales result in impaired alignment at border limits.

¹ The range between the spatial minimum and maximum estimated atmospheric deposition of sulfur and nitrogen for different locations in the state expressed in eq ha⁻¹ y⁻¹.

² One standard deviation of the weathering rate estimated for different locations in the state expressed as eq ha⁻¹ y⁻¹ of base cations released.

- Research has shown that sensitive forest areas show losses in forest health and productivity, and are more susceptible to climatic stress events, pests and diseases. For instance, preliminary results from the 30 sites of the Quebec Forest Monitoring Network (RESEF) show that forest sites in sensitive areas are growing 30% more slowly than sites located in tolerant areas. These findings point out the need to a more precise assessment of the risks associated with critical load exceedances.

Data variability within the NEG/ECP Region

The Forest Mapping Group recognized early on that the type and availability of the data necessary to perform critical load mapping for the region was not consistently uniform among the jurisdictions; consequently a certain amount of variability becomes evident upon close examination of the mapping results, particularly along the U.S./Canadian border. The Forest Mapping Workgroup agrees that each jurisdiction has been mapped in accordance with the Forest Mapping Protocol. The variability of the mapping results in no way detracts from or diminishes the impact or value of the data presented here as a tool with which to inform policymakers within the region. Some of the variability noted at the US/Canadian border is also due to differences in land use and landcover differences which can be pronounced in some border regions.

For example, the atmospheric deposition rates for sulphur and nitrogen which were used to identify exceedances of the calculated critical load values for forest ecosystems in the New England states are based on 1999-2003 total atmospheric deposition using a high-resolution deposition model (Miller, 2000; Miller et al., 2005); while they are based on the 1999-2002 total atmospheric deposition for forest ecosystems in Quebec and the Atlantic Canadian provinces (Shaw et al., 2006).

The critical load map for the Atlantic Provinces doesn't differentiate between agricultural and forested lands; whereas the maps for Quebec and the New England States include only forested land.

The variability in data attributes resulted in the New England states, the Atlantic provinces, and Quebec being mapped at different resolutions, varying from 0.09 ha grid cells for the New England states up to an average polygon size of 150 ha in Atlantic Canada (average polygon size in Quebec: 14.6 ha).

The soil mineral weathering rate estimation methods applied were also different among jurisdictions as permitted in the protocol (NEG/ECP Environment Task Group 2001). For the New England states, it was possible to use the data-intensive PROFILE model to estimate soil chemical weathering rates, while in Atlantic Canada the simpler soil substrate – texture model was used and its values seemed to correctly reflect forest soil conditions in Atlantic Canada (Whitfield et al., 2006). For Quebec, the latter model was applied with the use of a correction factor derived from the PROFILE model outputs from Maine used as calibration. The Forest mapping Group suggests that the evaluation of soil mineral weathering rates could be improved – and uncertainty reduced – through additional field studies.

Conclusions

More than one third of the mapped area of the NEG/ECP has been designated as sensitive to the extent that these areas are subjected to acid deposition in excess of their terrestrial critical loads. The possibility that high critical load exceedances leading directly or indirectly to poor forest growth and health in the exceeded regions appears strong. The critical load maps produced thus far demonstrate that further reductions in national and international S and N emission rates should be undertaken in order to protect forests from excessive soil acidification.

Additional work is necessary to improve upon the accuracy which could enhance and the utility of the regional critical load maps. Data variability between jurisdictions compromises our ability to draw seamless maps and a workshop bringing together forest mapping specialists and related disciplines in the region would build upon the success demonstrated to date.

Recommendations

1. In general, that a sustained incremental effort be made to improve the consistency of the data and the spatial resolution of the regional critical load map.
2. A regional forest sensitivity mapping workshop be organized within the next year.
3. An effort be made to improve the resolution of the soil weathering maps for the New England states, Quebec, and the Atlantic provinces.
4. Areas designated sensitive undergo further investigation to more precisely determine the scope and magnitude of the nutrient depletion occurring under current conditions of use and deposition.

References and deliverables

- Aherne, J., Ryan, D., de Kluizenaar, Y., van den Beuken, R. and Farrell, E.P., 2000. Literature review of the current state of knowledge regarding the calculation and mapping of critical loads: Determination and mapping of critical loads for sulphur and nitrogen and critical levels for ozone in Ireland, Environmental Protection Agency, Dublin.
- Arp, P.A., Leger, W., Moayeri, M.H. and Hurley, J.E., 2001. Methods for mapping Forest Sensitivity to Acid Deposition for northeastern North America. *Ecosystem Health*, 7: 35-47.
- DeMerchant, I., Hurley, J.E., Ouimet, R., Watmough, S.A. and Arp, P.A., 2004. Critical soil acidification loads and exceedances for Eastern Canada, Workshop report to Environment Canada, Fredericton, N.B.
- Jeffries, D.S. and Ouimet, R., 2005. Critical loads: are they being exceeded? Chapter 8. In: Environment Canada Science Assessment and Integration Branch (Editor),

- 2004 Scientific Assessment of Acid Deposition, Downsview, ON, Canada, pp. 341-369.
- Jeffries, D.S. et al., 2006. Combining aquatic and forest soil critical load and exceedance estimates for southeastern Canada. *Water Air and Soil Pollut. Focus*: (submitted at the 2005 Acid Rain Congress).
- Miller, E.K. 2000. Atmospheric Deposition to Complex Landscapes: HRDM – A Strategy for Coupling Deposition Models to a High-Resolution GIS. Proceedings of the National Atmospheric Deposition Program Technical Committee Meeting, October 17-20, 2000, Saratoga Springs, New York.
- Miller, E.K., VanArsdale, A., Keeler, G.J., Chalmers, A., Poissant, L., Kamman, N., and Brulotte, R. (2005) Estimation and Mapping of Wet and Dry Mercury Deposition across Northeastern North America. *Ecotoxicology* 14, 53-70.
- Miller, E.K., 2005. Assessment of forest sensitivity to nitrogen and sulfur deposition in New Hampshire and Vermont, Conference of the New England Governors and Eastern Canadian Premiers, Forest Mapping Group, Halifax, N.S.
- Miller, E.K., 2006. Assessment of forest sensitivity to nitrogen and sulfur deposition in Maine. Report prepared for the Maine Department of Environmental Protection, Bureau of Air Quality, Ecosystem Research Group Ltd., Norwich, VT.
- Moayeri, M., Meng, F., Rui, Arp, P.A. and Foster, N.W., 2001. Evaluating critical soil acidification loads and exceedances for a deciduous forest at the Turkey Lakes watershed. *Ecosystems*, 4: 555-567.
- NEG/ECP Environment Task Group, 2001. Critical load of sulphur and nitrogen assessment and mapping protocol for upland forests., NewEngland Governors and eastern canadian Premiers, Acid Rain Action Plan, Halifax, Canada.
- Nilsson, J. and Grennfelt, P., 1988. Critical loads for sulphur and nitrogen. Report from a workshop held at Skokloster, Sweden, Miljorapport 1988:15, Nordic Council of Ministers, Copenhagen, Denmark.
- Ouimet, R., Arp, P.A., Watmough, S.A., Aherne, J. and DeMerchant, I., 2006. Determination and mapping critical loads of acidity and exceedances for upland forest soils in Eastern Canada. *Water Air Soil Pollut.*, 172: 57-66.
- Ouimet, R., Duchesne, L., Houle, D. and Arp, P.A., 2001. Critical loads of atmospheric S and N deposition and current exceedances for Northern temperate and boreal forests in Quebec. *Water Air Soil Pollut. Focus*, 1(1/2): 119-134.
- Pardo, L.H., Robin, Abbott, M., Duarte, N. and Miller, E.K., 2005. Tree chemistry database (version 1.0). Gen. Techn. Rep. NE-324, USDA For. Serv., Northeast. Res. Stn., Delaware, OH.
- Shaw, M., Vet, R.J. and Zhang, L., 2006. Analyzed data fields from the National Atmospheric Chemistry Database (NAtChem) and Analysis Facility, Air Quality Research Division, Meteorological Service of Canada, Environment Canada, 4905 Dufferin St., Toronto, Ontario, Canada M3H 5T4.
- Watmough, S.A., Aherne, J., Arp, P.A., DeMerchant, I. and Ouimet, R., 2004. Canadian Experiences in Development of Critical Loads for Sulfur and Nitrogen. In: Aguirre, C. Bravo and et. al. (Editors), *Monitoring Science and Technology Symposium: Unifying Knowledge for Sustainability in the Western Hemisphere*. Proceedings RMRS-P-000. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Denver, CO.

Whitfield, C.J., Watmough, S.A., Aherne, J. and Dillon, P.J., 2006. A comparison of weathering rates for acid-sensitive catchments in Nova Scotia, Canada and their impact on critical load calculations. *Geoderma*, 136: 899 - 911.

The Forest Mapping Group

Paul A. Arp

Faculty of Forestry & Environmental Management
University of New Brunswick
28, Dineen Drive
Fredericton, N.B. E3B 6C2
Email: arp2@unb.ca

Eric Miller

President and Senior Scientist
Ecosystems Research Group, Ltd.
Aldrich House, Suite 11, 16 Beaver Meadow Rd.
PO Box 1227
Norwich, VT 05055
Tel: 802-649-5550
Fax: 802-649-5551
Email: ekmiller@ecosystems-research.com

Linda Pardo

USDA Forest Service
Northeastern Research Station
P.O. Box 968
Burlington, VT 05402
Tel: 802-951-6771, ext. 1330
Fax: 802-951-6368
Email: lpardo@fs.fed.us

Julian Aherne

Environmental and Resources Studies
Trent University
1600 West Bank Drive
Peterborough, ON K9J 7B8
Email: jaherne@trentu.ca

Sandy Wilmot

(member up to 2004)
Vermont Dept. of Forests, Parks & Recreation
Vermont Forest Ecosystem Monitoring
111 West Street
Essex Jct., VT 05452
Tel: 802-879-5687
Fax: 802-878-5192
Email: sandy.wilmot@anrmail.anr.state.vt.us

Scott Payne

Forestry Branch
Department of Forest Resources and Agrifoods
P.O. Box 2006
Fortis Building
Corner Brook, NF A2H 6J8
Tel: 709-637-2353
Fax: 709-637-2290
Email: sepayne@mail.gov.nf.ca

Ian DeMerchant

Canadian Forest Service
Atlantic Forestry Centre
P.O. Box 4000
Fredericton, N.B. E3B 5P7
Email: IDeMerch@NRCan.gc.ca

Rock Ouimet

Canadian co-chair
Direction de la recherche forestière
Ministère des Ressources Naturelles du Québec
2700, rue Einstein
Québec (QUÉBEC) G1P 3W8
Tel: 418-643-7994 (6533)
Fax: 418-643-2165
Email: rock.ouimet@mrnf.gouv.qc.ca

Shaun Watmough

Environmental and Resources Studies
Trent University
1600 West Bank Drive
Peterborough, ON K9J 7B8
Email: swatmough@trentu.ca

Wade Bowers

Canadian Forest Service
P.O. Box 960
Corner Brook, NF A2H 6J3
Tel: 709-637-4920
Fax: 709-637-4910
Email: wbowers@nrca.gc.ca

Ivan Downton

Director Forest Ecosystem Management
Forestry Branch
Department of Forest Resources and Agrifoods
P.O. Box 2006, Fortis Building
Corner Brook, NF A2H 6J8
Tel: 709-637-2284
Fax: 709-634-4378
Email: idownton@mail.gov.nf.ca

Joan Carlson

U.S. Dept. of Agriculture Forest Service
719 North Main Street
Laconia, NH 03246
Tel: (603) 528-8721
Fax: (603) 528-8783
Email: jcarlson/r9_whitemtn@fs.fed.us

Guy Fenech

(up to 2004)
Senior Science Advisor
Atmospheric Environment Service
Attention: APAC
Environment Canada
4905 Dufferin Street
Toronto, ON M3H 5T4
Tel: 416-739-4649
Fax: 416-739-4882
Email: guy.fenech@ec.gc.ca

Jennifer Bofinger

NH Division of Forests and Lands
172 Pembroke Road
P.O. Box 1856
Concord, NH 03302-1856
Tel: 603-271-7858
Fax: 603-271-2629
Email: j_bofinger@dred.state.nh.us

David Struble

Maine Forest Service
State House Station 22
Augusta, ME 04333
Tel: 207-287-4981
Fax: 207-287-8422
Email: dave.struble@state.me.us

Luc Lamontagne

Agriculture et Agroalimentaire Canada
2560, boul. Hochelaga
Sainte-Foy (Québec) G1V 2J3
Tel: 418-648-7730
Fax: 418-648-5489
Email: lamontagnel@em.agr.ca

Heather A. Morrison

Senior Science Advisor
Scientific Assessment and Integration Branch
Environment Canada
4905 Dufferin Street,
Downsview, ON,
M3H 5T4

Harry Hirvonen

(up to 2005)
Science Advisor, Forest Health
Science Branch, Canadian Forest Service
Natural Resources Canada
580 Booth St.
Ottawa, ON K1A 0E4
Tel: (613) 947-9015
Fax: (613) 947-9090
Email: hirvonen@nrcan.gc.ca

Wendy Leger

(up to 2003)
Water Projects Officer
Water Issues Division
Atmospheric Environment Branch
Environment Canada - Ontario Region
867 Lakeshore Road
Burlington, ON L7R 4A6
Tel: 905-336-4949
Fax: 905-336-8901
Email: wendy.leger@ec.gc.ca

Charles Burnham

Mass. Dept. of Environment
Division of Forest and Parks
100 Cambridge St., 19th floor
Boston, MA 02202
Tel: 617-727-3180, ext. 675
Fax: 617-727-9402
Email: charlie.burnham@state.ma.us

Cathy Sparks

Division of Forest Environment
1037 Hartford Pike
N. Scituate, RI 02857
Tel: 401-646-3367
Fax: 401-647-3590
Email: "Cathy Sparks"riforestry@edgenet.net

copy: Bob Dick *

Manager Forest Management Planning
NB Dept. of Natural Resources & Energy
P.O. Box 6000
Fredericton, NB E3B 5H1
Tel: 506-453-2516
Fax: 506-453-6689
Email: robert.dick@gov.nb.ca

** Does not participate in meetings or conference calls*

Silvina Carou

Air Quality Science Officer
Scientific Assessment and Integration Branch
Environment Canada
4905 Dufferin Street,
Downsview, ON,
M3H 5T4